



Fingolimod Dr.Reddy's

Pregnancy-Specific Patient Reminder Card

▼ This medicinal product is subject to additional monitoring in Australia. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. You can report side effects to your doctor, or directly at www.tga.gov.au/reporting-problems



1. Before starting Fingolimod Dr.Reddy's (fingolimod) treatment

Fingolimod Dr.Reddy's is contraindicated in pregnant women and women of child-bearing potential (including adolescents) not using effective contraception.

At treatment start and then regularly, your doctor will inform you about the teratogenic risk (causes defects to unborn babies) and required actions to minimise this risk.



A pregnancy test must be conducted and the negative result verified by a doctor before starting treatment.

Your doctor will inform you about the need for effective contraception while on treatment and for 2 months after discontinuation. Talk to your doctor about the most effective contraception options available to you.

Please read the Fingolimod Dr.Reddy's Patient's, Parents and Caregiver's Guide Leaflet provided by your doctor.



2. During the treatment

Women must not become pregnant during treatment and for 2 months after discontinuing treatment. Patients must use effective contraception while taking Fingolimod Dr.Reddy's.

Pregnancy tests must be repeated at suitable intervals. Women must not become pregnant during treatment and for 2 months after discontinuing treatment.

Pregnancy tests must be repeated at suitable intervals.



Your doctor will provide regular counselling about Fingolimod Dr.Reddy's serious risks to the foetus.

If you become pregnant or if you want to become pregnant please discuss this with your doctor because Fingolimod treatment must be discontinued.

In the event of a pregnancy your doctor will provide counselling.

Your doctor will give you medical advice regarding the harmful effects of Fingolimod Dr.Reddy's to the foetus and will provide an evaluation of the potential outcome.



3. After discontinuation

Inform your doctor immediately if you believe your MS is getting worse (e.g. weakness or visual changes) or if you notice any new symptoms after stopping treatment with Fingolimod due to pregnancy.

Effective contraception is needed for 2 months after stopping Fingolimod treatment because of the length of time it takes for Fingolimod to leave the body.



4. Reporting of Adverse Events

The safe use of Fingolimod Dr.Reddy's is of paramount importance. As part of the ongoing safety monitoring, Dr. Reddy's Laboratories wishes to learn of Adverse Events that have occurred during the use of Fingolimod.

For reporting an adverse events please contact Dr. Reddy's Laboratories,

DR. REDDYS' LABORATORIES CONTACT DETAILS

Tel: 1800 733

Email: ds@commercialeyes.com.au

Please report suspected adverse drug reactions (ADRs) to the Therapeutic Goods Administration (TGA) website www.tga.gov.au/reporting-problems.



Dr Reddy's Laboratories (Australia) Pty Ltd
Suite 3.03, Level 3, 390 St Kilda Road
Melbourne VIC 3004, Australia
©2025 March 2025

